

KAYNAKLAR

- Ağaoğlu YS. (1986). Üzümsü Meyveler. Ankara Üniversitesi Ziraat Fakültesi Yayınları No: 984 Ders Kitabı No: 290, 377s.
- Bothmer RV., Jacobsen N., Baden C., Jbrgensen RB., Linde-Laursen I. (1991). An ecogeographical study of the genus *Hordeum*. Systematic and Ecogeographic Studies on Crop Gene-pools 7, IBPGR, Rome, 126 pp.
- Brouat C., Gielly L., Mckey D. (2001). Phylogenetic relationships in the genus *Leonardoxa* (Leguminosae: Caesalpinioideae) inferred from chloroplast trnL intron and trnL-trnF intergenic spacer sequences. American Journal of Botany, 88: 143-149.
- Eck RV., Dayhoff MO. (1966). Atlas of protein sequence and structure. National Biomedical Research Foundation, Silver Spring.
- Eriksson T., Hibbs MS., Donoghues MJ. (1998). Phylogenetic analysis of *Potentilla* using DNA sequences of nuclear ribosomal internal transcribed spacers (ITS), and implications for the classification of Rosoideae (Rosaceae). Plant Systematics and Evolution, 211: 155-179.
- Eriksson T., Hibbs MS., Yoder AD., Delwiche CF., Donoghues MJ. (2003). The Phylogeny of Rosoideae (Rosaceae) based on sequences of the internal transcribed spacers (ITS) of nuclear ribosomal DNA and the trnL/F region of chloroplast DNA. International Journal of Plant Sciences, 164(2): 197-211.
- Fitch W. (1977). On the problem of discovering the most parsimonious tree. American Naturalist, 111: 223-257.
- Hall TA. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. Nucleic Acids Symposium Series, 41: 95-98.
- Haymes KM. (1996). Mini-prep method suitable for a plant breeding program. Plant Molecular Biology Reporter, 14 (3): 280-284.
- Karakoç D. (2011). Orta ve Doğu Karadeniz Bölgesi Doğal Florasındaki Böğürtlen Genotipleri Arasındaki Biyoçeşitliliğin Moleküler Belirteçlerle Saptanması. Gaziosmanpaşa Üniversitesi Fen Bilimleri Enstitüsü Yüksek Lisans Tezi, Tokat.
- Longhi S., Giongo L., Buti M., Surbanovski N., Viola R., Velasco R., Ward JA., Sargent DJ. (2014). Molecular genetics and genomics of the Rosoideae: state of the art and future perspectives. Horticulture Research, 1, 2052-7276/14.
- Onur C. (2006). Üzümsü meyveler ıslah programından sempozyumlara. II. Ulusal Üzümsü Meyveler Sempozyumu, 14-16 Eylül 2006, Tokat.
- Potter D., Eriksson T., Evans RC., Oh S., Smedmark JEE., Morgan DR., Kerr M., Robertson KR., Arsenault M., Dickinson TA., Campbell, CS. (2007). Phylogeny and classification of Rosaceae. Plant Systematics and Evolution, 266: 5-43.
- Saitou N., Nei M. (1987). The Neighbor-Joining method: a new method for reconstructing phylogenetic trees. Molecular Biology Evolution, 4: 406-425.
- Sochor M., Vasut RJ., Sharbel TF., Travnicek B. (2015). How just a few makes a lot: speciation via reticulation and *Apomixis* on example of European Brambles (*Rubus* subgen. *Rubus*, Rosaceae). Molecular Phylogenetics and Evolution, 89: 13-27.
- Soejima A., Nagamasu H. (2004). Phylogenetic analysis of Asian *Symplocos* (Symplocaceae) based on nuclear and chloroplast DNA sequences. Journal of Plant Research, 117(3):199-207.
- Tamura K., Stecher G., Peterson, D. Filipski A., Kumar S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. Molecular Biology Evolution, 30: 2725-2729.
- Thompson JD., Gibson TJ., Plewniak F., Jeanmougin F., Higgins DG. (1997). The clustal x-windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Research, 25: 4876-4882.

©Türkiye Herboloji Derneği, 2018

Geliş Tarihi/ Received: Ağustos/Agust, 2018
Kabul Tarihi/ Accepted: Kasım/November, 2018

To Cite: Koloren O. and Eker S. (2018). Molecular Diagnosis and Phylogeny of the *Rubus* Species of Ordu Province (In Turkish with English Abstract). Turk J Weed Sci, 21(2):39-45.

Alıntı için: Koloren O. ve Eker S. (2018). Ordu İli *Rubus* Türlerinin Moleküler Tanısı ve Filogenisi. Turk J Weed Sci, 21(2):39-45.